

# Drug Policy:

## Iwilfin™ (eflornithine)

|   |   |   |   |
|---|---|---|---|
| <b>POLICY NUMBER</b><br>UM ONC_1495                   | <b>SUBJECT</b><br>Iwilfin™ (eflornithine) | <b>DEPT/PROGRAM</b><br>UM Dept  | <b>PAGE 1 OF 3</b>                                    |
| <b>DATES COMMITTEE REVIEWED</b><br>01/10/24, 02/14/24 | <b>APPROVAL DATE</b><br>February 14, 2024 | <b>EFFECTIVE DATE</b><br>February 23, 2024                            | <b>COMMITTEE APPROVAL DATES</b><br>01/10/24, 02/14/24 |
| <b>PRIMARY BUSINESS OWNER: UM</b>                     |   | <b>COMMITTEE/BOARD APPROVAL</b><br>Utilization Management Committee   |   |
| <b>NCQA STANDARDS</b><br>UM 2                         |   | <b>ADDITIONAL AREAS OF IMPACT</b>                                     |   |
| <b>CMS REQUIREMENTS</b>                               | <b>STATE/FEDERAL REQUIREMENTS</b>         | <b>APPLICABLE LINES OF BUSINESS</b><br>Commercial, Exchange, Medicaid |   |

### I. PURPOSE

To define and describe the accepted indications for Iwilfin (eflornithine) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

Evolent is responsible for processing all medication requests from network ordering providers. Medications not authorized by Evolent may be deemed as not approvable and therefore not reimbursable.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

### II. INDICATIONS FOR USE/INCLUSION CRITERIA

#### A. Continuation requests for a not-approvable medication shall be exempt from this Evolent policy provided:

1. The member has not experienced disease progression on the requested medication **AND**
2. The requested medication was used within the last year without a lapse of more than 30 days of having an active authorization **AND**
3. Additional medication(s) are not being added to the continuation request.

#### B. Neuroblastoma

1. Iwilfin (eflornithine) may be used as monotherapy to reduce the risk of relapse in adult and pediatric members with high-risk neuroblastoma (HRNB) who have demonstrated at least a partial response to prior multiagent, multimodality therapy including anti-GD2 immunotherapy (i.e. dinutuximab, naxitamab).
  - a. Iwilfin (eflornithine) may be used until disease progression, unacceptable toxicity, or for a maximum of 2 years.

### III. EXCLUSION CRITERIA

- A. Disease progression while taking Iwifin (eflornithine).
- B. Concurrent use with other anticancer therapies.
- C. Dosing exceeds single dose limit of Iwifin (eflornithine) 768 mg.
- D. Investigational use of Iwifin (eflornithine) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
  - 1. Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
  - 2. Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
  - 3. Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definitions of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of < 0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.
  - 4. Whether the experimental design, considering the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
  - 5. That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
  - 6. That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
  - 7. That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

### IV. MEDICATION MANAGEMENT

- A. Please refer to the FDA label/package insert for details regarding these topics.

### V. APPROVAL AUTHORITY

- A. Review – Utilization Management Department
- B. Final Approval – Utilization Management Committee

### VI. ATTACHMENTS

- A. None

### VII. REFERENCES

- A. Iwifin prescribing information 2023. USWM, LLC Louisville, KY 40241
- B. Sholler, G.L.S., Ferguson, W., Bergendahl, G. et al. Maintenance DFMO Increases Survival in High Risk Neuroblastoma. *Sci Rep* **8**, 14445 (2018). <https://doi.org/10.1038/s41598-018-32659-w>

- C. Oesterheld, J, et al. Eflornithine as Post immunotherapy Maintenance in High-Risk Neuroblastoma: Externally Controlled, Propensity Score-Matched Survival Outcome Comparisons. Journal of Clinical Oncology. DOI: [10.1200/JCO.22.0287](https://doi.org/10.1200/JCO.22.0287)
- D. Clinical Pharmacology Elsevier Gold Standard 2023.
- E. Micromedex® Healthcare Series: Micromedex Drugdex Ann Arbor, Michigan 2023.
- F. National Comprehensive Cancer Network. Cancer Guidelines and Drugs and Biologics Compendium 2023.
- G. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. J Clin Oncol. 2014 Apr 20;32(12):1277-80.
- H. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>.