

Drug Policy:

Treanda/Bendeka/Belrapzo/Vivimusta™ (bendamustine)

POLICY NUMBER UM ONC_1215	SUBJECT Treanda/Bendeka/Belrapzo/Vivimusta™ (bendamustine)		DEPT/PROGRAM UM Dept	PAGE 1 of 3
DATES COMMITTEE REVIEWED 09/12/12, 02/12/14, 12/17/15, 12/14/17, 03/14/18, 03/07/19, 04/10/19, 12/11/19, 04/08/20, 01/13/21, 11/15/21, 12/08/21, 03/09/22, 05/11/22, 09/14/22, 03/08/23, 05/10/23, 09/13/23, 08/14/24	APPROVAL DATE August 14, 2024	EFFECTIVE DATE August 30, 2024	COMMITTEE APPROVAL DATES 09/12/12, 02/12/14, 12/17/15, 12/14/17, 03/14/18, 03/07/19, 04/10/19, 12/11/19, 04/08/20, 01/13/21, 11/15/21, 12/08/21, 03/09/22, 05/11/22, 09/14/22, 03/08/23, 05/10/23, 09/13/23, 08/14/24	
PRIMARY BUSINESS OWNER: UM		COMMITTEE/BOARD APPROVAL Utilization Management Committee		
NCQA STANDARDS UM 2		ADDITIONAL AREAS OF IMPACT		
CMS REQUIREMENTS	STATE/FEDERAL REQUIREMENTS		APPLICABLE LINES OF BUSINESS Commercial, Exchange, Medicaid	

I. PURPOSE

To define and describe the accepted indications for Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

Evolent is responsible for processing all medication requests from network ordering providers. Medications not authorized by Evolent may be deemed as not approvable and therefore not reimbursable.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

II. INDICATIONS FOR USE/INCLUSION CRITERIA

A. Continuation requests for a not-approvable medication shall be exempt from this Evolent policy provided:

1. The requested medication was used within the last year, AND
2. The member has not experienced disease progression and/or no intolerance to the requested medication, AND
3. Additional medication(s) are not being added to the continuation request.

B. Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma

1. Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) may be used in combination with rituximab/rituximab biosimilar as initial or subsequent therapy for members with CLL.

C. Non-Hodgkin's Lymphoma

1. Indolent B-Cell Lymphomas: Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) may be used in the following clinical settings:
 - a. In combination with rituximab/rituximab biosimilar for primary or subsequent therapy of any of the following:
 - i. Follicular B-Cell Lymphoma
 - ii. Nodal Marginal Zone/Extra-Nodal Marginal Zone/Gastric MALT Lymphoma/Non-Gastric MALT Lymphoma/Splenic Marginal Zone Lymphoma.
2. Diffuse Large B-Cell Lymphoma
 - a. Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) may be used as subsequent therapy for relapsed or refractory disease in combination with Polivy (polatuzumab) with or without rituximab/rituximab biosimilar.
3. Mantle Cell Lymphoma
 - a. Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) may be used as initial or subsequent therapy in combination with rituximab/rituximab biosimilar.

D. Small Cell Lung Cancer (SCLC)

1. NOTE: Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) products are not supported by Evolent Policy for the treatment of relapsed/refractory SCLC. This policy position is based on the lack of Level 1 Evidence (randomized clinical trials and/or meta-analyses) to show superior outcomes compared to Evolent recommended regimens, available at <https://pathways.newcenturyhealth.com>.

III. EXCLUSION CRITERIA

- A. Member has disease progression while on Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine).
- B. Dosing exceeds single dose limit of Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) 100 mg/m² for CLL and 120 mg/m² for NHL.
- C. Treatment with Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) exceeds the maximum duration limit of 8 cycles for NHL and 6 cycles for CLL.
- D. Investigational use of Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
 1. Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
 2. Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
 3. Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definitions of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of less than 0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.

4. Whether the experimental design, considering the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
5. That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
6. That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
7. That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

IV. MEDICATION MANAGEMENT

- A. Please refer to the FDA label/package insert for details regarding these topics.

V. APPROVAL AUTHORITY

- A. Review – Utilization Management Department
- B. Final Approval – Utilization Management Committee

VI. ATTACHMENTS

- A. None

VII. REFERENCES

- A. Treanda prescribing information. Cephalon, Inc. Frazer, PA. 2021.
- B. Vivimusta prescribing information. Slayback Pharma LLC Princeton, NJ 2023
- C. Bendeka prescribing information. Teva Pharmaceuticals USA, Inc. North Wales, PA 2021,
- D. Belrapzo prescribing information. Eagle Pharmaceuticals, Inc. Woodcliff Lake, NJ. 2021.
- E. Clinical Pharmacology Elsevier Gold Standard 2023.
- F. Micromedex® Healthcare Series: Micromedex Drugdex Ann Arbor, Michigan 2023.
- G. National Comprehensive Cancer Network. Cancer Guidelines and Drugs and Biologics Compendium 2023.
- H. AHFS Drug Information. American Society of Health-Systems Pharmacists or Wolters Kluwer Lexi-Drugs. Bethesda, MD 2023.
- I. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. J Clin Oncol. 2014 Apr 20;32(12):1277-80.
- J. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>.