

Drug Policy:

Columvi™ (glofitamab-gxbm)

| | | | | |
|-------------------------------------------------------|----------------------------------------------|---------------------------------------------------------------------|-----------------------------------------------------------------------|--------------------|
| POLICY NUMBER UM ONC_1480 | SUBJECT Columvi™ (glofitamab-gxbm) | | DEPT/PROGRAM UM Dept | PAGE 1 OF 3 |
| DATES COMMITTEE REVIEWED 08/09/23, 08/14/24 | APPROVAL DATE August 14, 2024 | EFFECTIVE DATE August 30, 2024 | COMMITTEE APPROVAL DATES 08/09/23, 08/14/24 | |
| PRIMARY BUSINESS OWNER: UM | | COMMITTEE/BOARD APPROVAL Utilization Management Committee | | |
| NCQA STANDARDS UM 2 | | ADDITIONAL AREAS OF IMPACT | | |
| CMS REQUIREMENTS | STATE/FEDERAL REQUIREMENTS | | APPLICABLE LINES OF BUSINESS Commercial, Exchange, Medicaid | |

I. PURPOSE

To define and describe the accepted indications for Columvi (glofitamab-gxbm) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

Evolent is responsible for processing all medication requests from network ordering providers. Medications not authorized by Evolent may be deemed as not approvable and therefore not reimbursable.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

II. INDICATIONS FOR USE/INCLUSION CRITERIA

A. Continuation requests for a not-approvable medication shall be exempt from this Evolent policy provided:

1. The member has not experienced disease progression on the requested medication **AND**
2. The requested medication was used within the last year without a lapse of more than 30 days of having an active authorization **AND**
3. Additional medication(s) are not being added to the continuation request.

B. Diffuse Large B-Cell Lymphoma (DLBCL)

1. Columvi (glofitamab-gxbm) may be used as monotherapy for relapsed or refractory* DLBCL or large B-cell lymphoma (LBCL) arising from follicular lymphoma, **AND** the member has failed at least two systemic therapies, including an anti-CD20 monoclonal antibody (e.g., R-CHOP, R-ICE, R-DHAP, R-ESHAP, R-EPOCH, R-GDP).

*includes patients with disease progression after transplant or CAR T-cell therapy

III. EXCLUSION CRITERIA

- A. Disease progression while taking Columvi (glofitamab-gxbm)
- B. Concurrent use with other anticancer therapies.
- C. Dosing exceeds single dose limit of Columvi (glofitamab-gxbm) 30 mg.
- D. Investigational use of Columvi (glofitamab-gxbm) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
 - 1. Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
 - 2. Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
 - 3. Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definitions of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of less than 0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.
 - 4. Whether the experimental design, considering the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
 - 5. That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
 - 6. That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
 - 7. That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

IV. MEDICATION MANAGEMENT

- A. Please refer to the FDA label/package insert for details regarding these topics.

V. APPROVAL AUTHORITY

- A. Review – Utilization Management Department
- B. Final Approval – Utilization Management Committee

VI. ATTACHMENTS

- A. None

VII. REFERENCES

- A. Dickinson et al. Glofitamab for relapsed or refractory Diffuse Large B-Cell Lymphoma. December 15, 2022. N Engl J Med 2022; 387:2220-2231. DOI: 10.1056/NEJMoa2206913
- B. Columvi prescribing information. Genentech USA, Inc. South San Francisco, CA 94080. 2023

- C. AHFS Drug Information. American Society of Health-Systems Pharmacists or Wolters Kluwer Lexi-Drugs Bethesda, MD 2024.
- D. Clinical Pharmacology Elsevier Gold Standard 2024.
- E. Micromedex® Healthcare Series: Micromedex Drugdex Ann Arbor, Michigan 2024.
- F. National Comprehensive Cancer Network. Cancer Guidelines and Drugs and Biologics Compendium 2024.
- G. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. J Clin Oncol. 2014 Apr 20;32(12):1277-80.
- H. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services:
<https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>.