

Reference number(s) 1649-A

Specialty Guideline Management bosentan-Tracleer

Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over-the-counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Tracleer	bosentan

Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-approved Indications^{1,2}

Indicated for the treatment of pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1):

- In adults to improve exercise ability and to decrease clinical worsening. Studies establishing
 effectiveness included predominantly patients with WHO Functional Class II-IV symptoms and
 etiologies of idiopathic or heritable PAH, PAH associated with connective tissue diseases, and
 PAH associated with congenital heart disease with left-to-right shunts.
- In pediatric patients aged 3 years and older with idiopathic or congenital PAH to improve pulmonary vascular resistance (PVR), which is expected to result in an improvement in exercise ability.

All other indications are considered experimental/investigational and not medically necessary.

bosentan-Tracleer SGM 1649-A P2024a_R.docx

© 2024 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

Prescriber Specialties

This medication must be prescribed by or in consultation with a pulmonologist or cardiologist.

Coverage Criteria

Pulmonary Arterial Hypertension (PAH)¹⁻⁶

Authorization of 12 months may be granted for treatment of PAH when ALL of the following criteria are met:

- Member has PAH defined as WHO Group 1 class of pulmonary hypertension (refer to Appendix).
- PAH was confirmed by either of the following criteria:
 - Pretreatment right heart catheterization with all of the following results:
 - Mean pulmonary arterial pressure (mPAP) > 20 mmHg
 - Pulmonary capillary wedge pressure (PCWP) ≤ 15 mmHg
 - Pulmonary vascular resistance (PVR) > 2 Wood units. For pediatric members, pulmonary vascular resistance index (PVRI) > 3 Wood units x m² is also acceptable.
 - For infants less than one year of age, PAH was confirmed by Doppler echocardiogram if right heart catheterization cannot be performed.

Continuation of Therapy

Authorization of 12 months may be granted for members with an indication listed in the coverage criteria section who are currently receiving the requested medication through a paid pharmacy or medical benefit, and who are experiencing benefit from therapy as evidenced by disease stability or disease improvement.

Appendix

WHO Classification of Pulmonary Hypertension (PH)⁴

Note: Patients with heritable PAH or PAH associated with drugs and toxins might be long-term responders to calcium channel blockers.

bosentan-Tracleer SGM 1649-A P2024a_R.docx

© 2024 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

Group 1: Pulmonary Arterial Hypertension (PAH)

- Idiopathic
 - Long-term responders to calcium channel blockers
- Heritable
- Associated with drugs and toxins
- Associated with:
 - Connective tissue disease
 - Human immunodeficiency virus (HIV) infection
 - Portal hypertension
 - Congenital heart disease
 - Schistosomiasis
- PAH with features of venous/capillary (pulmonary veno-occlusive disease [PVOD]/pulmonary capillary hemangiomatosis [PCH]) involvement
- Persistent PH of the newborn

Group 2: PH associated with Left Heart Disease

- Heart failure:
 - With preserved ejection fraction
 - With reduced or mildly reduced ejection fraction
 - Cardiomyopathies with specific etiologies (i.e., hypertrophic, amyloid, Fabry disease, and Chagas disease)
- Valvular heart disease:
 - Aortic valve disease
 - Mitral valve disease
 - Mixed valvular disease
- Congenital/acquired cardiovascular conditions leading to post-capillary PH

Group 3: PH associated with Lung Diseases and/or Hypoxia

- Chronic obstructive pulmonary disease (COPD) and/or emphysema
- Interstitial lung disease
- Combined pulmonary fibrosis and emphysema
- Other parenchymal lung diseases (i.e., parenchymal lung diseases not included in Group 5)
- Nonparenchymal restrictive diseases:
 - Hypoventilation syndromes
 - Pneumonectomy
- Hypoxia without lung disease (e.g., high altitude)
- Developmental lung diseases

Group 4: PH associated with Pulmonary Artery Obstructions

- Chronic thromboembolic PH
- Other pulmonary artery obstructions
 - Sarcomas (high- or intermediate-grade or angiosarcoma)

bosentan-Tracleer SGM 1649-A P2024a_R.docx

© 2024 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

- Other malignant tumors (e.g., renal carcinoma, uterine carcinoma, germ-cell tumors of the testis)
- Non-malignant tumors (e.g., uterine leiomyoma)
- Arteritis without connective tissue disease
- Congenital pulmonary artery stenoses
- Hydatidosis

Group 5: PH with Unclear and/or Multifactorial Mechanisms

- Hematological disorders, including inherited and acquired chronic hemolytic anemia and chronic myeloproliferative disorders
- Systemic disorders: Sarcoidosis, pulmonary Langerhans cell histiocytosis, and neurofibromatosis type 1
- Metabolic disorders, including glycogen storage diseases and Gaucher disease
- Chronic renal failure with or without hemodialysis
- Pulmonary tumor thrombotic microangiopathy
- Fibrosing mediastinitis
- · Complex congenital heart disease

References

- 1. Tracleer [package insert]. Titusville, NJ: Actelion Pharmaceuticals US, Inc.; February 2024.
- 2. Bosentan [package insert]. Parsippany, NJ: Teva Pharmaceuticals. March 2024.
- 3. Simonneau G, Montani D, Celermajer DS, et al. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur Respir J.* 2019;53(1):1801913. doi:10.1183/13993003.01913-2018
- 4. Kovacs G, Bartolome S, Denton CP, et al. Definition, classification and diagnosis of pulmonary hypertension. Eur Respir J. 2024;64(4):2401324. doi: 10.1183/13993003.01324-2024
- 5. Chin KM, Gaine SP, Gerges C, et al. Treatment algorithm for pulmonary arterial hypertension. Eur Respir J. 2024;64(4):2401325. doi: 10.1183/13993003.01325-2024
- 6. Ivy D, Rosenzweig EB, Abman SH, et al. Embracing the challenges of neonatal and paediatric pulmonary hypertension. Eur Respir J. 2024;64(4):2401345. doi: 10.1183/13993003.01345-2024